THE LONG-TAILED JAY.

BY JOSEPH GRINNELL.

Aphelocoma californica immanis, new subspecies.

Subsp. Char. —In coloration similar to Aphelocoma californica, but size greater and tail proportionately much longer.

Type. — \mathcal{J} ad.. No. 4582, coll. J. G.; Scio, Oregon, January 9, 1901; collected by A. G. Prill and procured for me by B. J. Bretherton.

Habitat. - Willamette Valley, Oregon (Scio and Salem).

MEASUREMENTS 1 OF SPECIMENS.

No., Coll. J. G.	Sex.	Date.	Locality.	Wing.	Tail.	Tarsus.	Culm.	Bill from Nostril.	Depth of Bill,
4585	8	March 30, '00	Scio, Oregon	127	151	44	27	20	10.5
4582	♂	Jan. 9, '01	66 66	133	165	44	28	21.5	10
4586	ç	Feb. 8, 'or	64 64	125	152	42	28	20.5	10
4583	9	March 3, '00		129	150	41	27	21	10

I In millimeters.

GENERAL NOTES.

Brünnich's Murre in Lewis and Jefferson Counties, New York.—On Nov. 29, 1900, I had an adult male specimen of *Uria lomvia* brought to me by a farmer who stated he caught him alive in a mud hole near his home. This is, I believe, the first instance of the capture of this species in Lewis County.

A few days later, Dec. 4, I received another from a gunner at Redwood, Jefferson County, who stated there was quite a large flock of them on the lake at that place. This specimen had eight small black bass in its stomach, the largest three inches long.—JAMES H. MILLER, Lowville, Lewis Co., N. Y.